(Contents continued from back cover)

Comparison between aqueous and non-aqueous solvent evaporation methods for microencapsulation of drug–resin complexes
D. Torres, L. Boado, D. Blanco, J.L. Vila-Jato (Spain) 171
Skin permeation model of phenyl alcohols: comparison of experimental conditions
A. López, V. Faus, O. Díez-Sales, M. Herráez (Spain) 183
EPR study of mucoadhesive ointments for delivery of liposomes into the oral mucosa
M. Petelin, M. Šentjurc, Z. Stolič, U. Skalerič (Slovenia) 193
Ex vivo oral mucosal permeation of lidocaine hydrochloride with sucrose fatty acid esters as absorption enhancers
A. Ganem-Quintanar, D. Quintanar-Guerrero, F. Falson-Rieg, P. Buri (Mexico, France, Switzerland) 203
A flow-through dissolution method for a two component drug formulation where the actives have markedly differing solubility properties
W.C.G. Butler, S.R. Bateman (UK) 211
Hydrolysis kinetics and stability predictions for a mixture of R and S temocillin isomers
S.C. Dyar, R.E. Notari (USA) 221
Tableting of controlled release multiparticulates, the effect of millisphere size and protective overcoating
J.L. Haslam, A.E. Forbes, G.S. Rork, T.L. Pipkin, D.A. Slade, D. Khossravi (USA) 233
Design of inhalation dry powder of pranlukast hydrate to improve dispersibility by the surface modification with light anhydrous silicic acid (AEROSIL 200)
Y. Kawashima, T. Serigano, T. Hino, H. Yamamoto, H. Takeuchi (Japan) 243

Book review 253

Short communication
The improved everted gut sac: a simple method to study intestinal P-glycoprotein
L. Barthe, M. Bessouet, J.F. Woodley, G. Houin (France, UK) 255

Noticeboard 259
Author index 263